TABLE 1. A SUMMARY (including footnotes) OF ATLANTIC MARINE MAMMAL STOCKASSESSMENT REPORTS FOR STOCKS OF MARINE MAMMALS UNDER NMFS AUTHORITYTHAT OCCUPY WATERS UNDER USA JURISDICTION.

Total Annual S.I. (serious injury) and Mortality and Annual Fisheries S.I. and Mortality are mean annual figures for the period 2011-2015. The "SAR revised" column indicates 2017 stock assessment reports that have been revised relative to the 2016 reports (Y=yes, N=no). If abundance, mortality, PBR or status have been revised, they are indicated with the letters "a", "m", "p" and "status" respectively. For those species not updated in this edition, the year of last revision is indicated. Unk = unknown and undet=undetermined (PBR for species with outdated abundance estimates is considered "undetermined").

| Species | Stock Area | NMFS Ctr. | Nbest | Nbest CV | Nmin | Rmax | Fr | PBR | r 1 | Annual Fish. S.I. and Mort. (cv) | Strategic Status | SAR Revised |
|-------------------------------|------------------------|-----------|-------|-------------|-------|-------------------|-----|-----|-------------------|-------------------------------------|---------------------|---------------------------|
| North Atlantic right whale | Western North Atlantic | NEC | 458 | 0 | 455 | 0.04 ^a | 0.1 | 0.9 | 5.36 ^a | 4.55 ^a | Y | Y (a, m, p) |
| Humpback whale | Gulf of Maine | NEC | 335 | 0.42 | 239 | 0.065 | 0.5 | 3.7 | 8.5 ^b | 6.7 ^b | Y | Y (a, m, p, status) |
| Fin whale | Western North Atlantic | NEC | 1,618 | 0.33 | 1,234 | 0.04 | 0.1 | 2.5 | 2.65° | 1.05° | Y | Y (m) |
| Sei whale | Nova Scotia | NEC | 357 | 0.52 | 236 | 0.04 | 0.1 | 0.5 | 0.8^{d} | 0 ^d | Y | N (2016) |
| Minke whale | Canadian east coast | NEC | 2,591 | 0.81 | 1,425 | 0.04 | 0.5 | 14 | 9.15 ^e | 7.75° | N | Y (m) |
| Blue whale | Western North Atlantic | NEC | unk | unk | 440 | 0.04 | 0.1 | 0.9 | unk | unk | Y | N (2010) |
| Sperm whale | North Atlantic | NEC | 2,288 | 0.28 | 1,815 | 0.04 | 0.1 | 3.6 | 0.8 | 0.8 | Y | N (2014) |

| Species | Stock Area | NMFS Ctr. | Nbest | Nbest CV | Nmin | Rmax | Fr | PBR | Total Annual S.I and Mort. | Annual Fish. S.I. and Mort. (cv) | Strategic Status | SAR Revised |
|------------------------------|------------------------|-----------|---------------------|--------------------|---------------------|------|-----|-----|-------------------------------|-------------------------------------|---------------------|----------------|
| Dwarf sperm whale | Western North Atlantic | SEC | 3,785 ^h | 0.47 ^k | 2,598 ^h | 0.04 | 0.4 | 21 | 3.5 | 3.5 (1.0) | N | N (2016) |
| Pygmy sperm whale | Western North Atlantic | SEC | 3,785 ^h | 0.47 ^k | 2,598 ^h | 0.04 | 0.4 | 21 | 3.5 | 3.5 (1.0) | N | N (2016) |
| Killer whale | Western North Atlantic | NEC | unk | unk | unk | 0.04 | 0.5 | unk | 0 | 0 | N | N (2014) |
| Pygmy killer whale | Western North Atlantic | SEC | unk | unk | unk | 0.04 | 0.5 | unk | 0 | 0 | N | N (2007) |
| False killer whale | Western North Atlantic | SEC | 442 | 1.06 | 212 | 0.04 | 0.5 | 2.1 | unk | unk | Y | N (2014) |
| Northern bottlenose whale | Western North Atlantic | NEC | unk | unk | unk | 0.04 | 0.5 | unk | 0 | 0 | N | N (2014) |
| Cuvier's beaked whale | Western North Atlantic | NEC | 6,532 | 0.32 | 5,021 | 0.04 | 0.5 | 50 | 0.4 | 0.2 | N | N (2013) |
| Blainville's beaked whale | Western North Atlantic | NEC | 7,092 ^g | 0.54 | 4,632 ^g | 0.04 | 0.5 | 46 | 0.2 | 0.2 | N | N (2013) |
| whale | Western North Atlantic | | 7,092 ^g | 0.54 | 4,632 ^g | 0.04 | 0.5 | 46 | 0 | 0 | N | N (2013) |
| Sowerby's beaked whale | Western North Atlantic | NEC | 7,092 ^g | 0.54 | 4,632 ^g | 0.04 | 0.5 | 46 | 0 | 0 | N | N (2014) |
| True's beaked whale | Western North Atlantic | NEC | 7,092 ^g | 0.54 | 4,632 ^g | 0.04 | 0.5 | 46 | 0 | 0 | N | N (2013) |

| Species | Stock Area | NMFS Ctr. | Nbest | Nbest CV | Nmin | Rmax | Fr | PBR | Total Annual S.I and Mort. | Annual Fish. S.I. and Mort. (cv) | Strategic Status | SAR Revised |
|----------------------------------|------------------------|-----------|--------|-------------|--------|------|------|-----|-------------------------------|-------------------------------------|---------------------|----------------|
| Melon-headed whale | Western North Atlantic | SEC | unk | unk | unk | 0.04 | 0.5 | unk | 0 | 0 | N | N (2007) |
| Risso's dolphin | Western North Atlantic | NEC | 18,250 | 0.46 | 12,619 | 0.04 | 0.48 | 126 | 43.2 | 43 (0.27) | N | Y (m) |
| Pilot whale, long- finned | Western North Atlantic | NEC | 5,636 | 0.63 | 3,464 | 0.04 | 0.5 | 35 | 38 | 38 (0.15) | Y | N (2016) |
| Pilot whale, short- finned | Western North Atlantic | SEC | 21,515 | 0.37 | 15,913 | 0.04 | 0.5 | 159 | 192 | 192 (0.17) | Y | N (2016) |
| Atlantic white- sided dolphin | Western North Atlantic | NEC | 48,819 | 0.61 | 30,403 | 0.04 | 0.5 | 304 | 57 | 57 (0.15) | N | Y (m) |
| White-beaked dolphin | Western North Atlantic | NEC | 2,003 | 0.94 | 1,023 | 0.04 | 0.5 | 10 | 0 | 0 | N | N (2007) |
| Common dolphin | Western North Atlantic | NEC | 70,184 | 0.28 | 55,690 | 0.04 | 0.5 | 557 | 437 | 437 (0.10) | N | Y (m) |
| Atlantic spotted dolphin | Western North Atlantic | | 44.715 | 0.43 | 31,610 | 0.04 | 0.5 | 316 | 0 | 0 | N | N (2013) |
| Pantropical spotted dolphin | Western North Atlantic | SEC | 3,333 | 0.91 | 1,733 | 0.04 | 0.5 | 17 | 0 | 0 | N | N (2013) |
| Striped dolphin | Western North Atlantic | NEC | 54,807 | 0.3 | 42,804 | 0.04 | 0.5 | 428 | 0 | 0 | N | N (2013) |
| Fraser's dolphin | Western North Atlantic | SEC | unk | unk | unk | 0.04 | 0.5 | unk | 0 | 0 | N | N (2007) |

| Species | Stock Area | NMFS Ctr. | Nbest | Nbest CV | Nmin | Rmax | Fr | PBR | Total Annual S.I and Mort. | Annual Fish. S.I. and Mort. (cv) | Strategic Status | SAR Revised |
|------------------------------|--|-----------|---------------------|-------------|---------------------|------|-----|-------|-------------------------------|-------------------------------------|---------------------|----------------|
| Rough-toothed dolphin | Western North Atlantic | SEC | 271 | 1.0 | 134 | 0.04 | 0.5 | 1.3 | 0 | 0 | N | N (2013) |
| Clymene dolphin | Western North Atlantic | SEC | unk | unk | unk | 0.04 | 0.5 | undet | 0 | 0 | N | N (2013) |
| Spinner dolphin | Western North Atlantic | SEC | unk | unk | unk | 0.04 | 0.5 | unk | 0 | 0 | Ν | N (2013) |
| Common bottlenose dolphin | Western North Atlantic, offshore | SEC | 77,532 ^f | 0.40 | 56,053 ^f | 0.04 | 0.5 | 561 | 39.4 | 39.4 (0.29) | N | N (2016) |
| Common bottlenose dolphin | Western North Atlantic, northern migratory coastal | SEC | 6,639 | 0.41 | 4,759 | 0.04 | 0.5 | 48 | 6.1-13.2 | 6.1-13.2 | Y | Y (a, m, p) |
| Common bottlenose dolphin | Western North Atlantic, southern migratory coastal | SEC | 3,751 | .060 | 2,353 | 0.04 | 0.5 | 23 | 0-14.3 | 0-14.3 | Y | Y (a, m, p) |
| Common bottlenose dolphin | Western North Atlantic, S. Carolina/Georgia coastal | SEC | 6,027 | 0.34 | 4,569 | 0.04 | 0.5 | 46 | 1.4-1.6 | 1.0-1.2 | Y | Y (a, m, p) |
| Common bottlenose dolphin | Western North Atlantic, northern Florida coastal | SEC | 877 | 0.49 | 595 | 0.04 | 0.5 | 6.0 | 0.6 | 0 | Y | Y (a, m, p) |
| Common bottlenose dolphin | Western North Atlantic, central Florida coastal | SEC | 1,218 | 035 | 913 | 0.04 | 0.5 | 9.1 | 0.4 | 0.4 | Y | Y (a, m, p) |
| Common bottlenose dolphin | Northern North Carolina Estuarine System | SEC | 823 | 0.06 | 782 | 0.04 | 0.5 | 7.8 | 0.8-18.2 | 0.2-17.6 | Y | Y (m) |

| Species | Stock Area | NMFS Ctr. | Nbest | Nbest CV | Nmin | Rmax | Fr | PBR | Total Annual S.I and Mort. | Annual Fish. S.I. and Mort. (cv) | Strategic Status | SAR Revised |
|------------------------------|---|-----------|--------|-------------|--------|-------|-----|-------|-------------------------------|-------------------------------------|---------------------|----------------|
| Common bottlenose dolphin | Southern North Carolina Estuarine System | SEC | unk | unk | unk | 0.04 | 0.5 | undet | 0.4-0.6 | 0.4-0.6 | Y | Y (m) |
| Common bottlenose dolphin | Northern South Carolina Estuarine System | SEC | unk | unk | unk | 0.04 | 0.5 | unk | 0.2 | 0.2 | Y | N (2015) |
| Common bottlenose dolphin | Charleston Estuarine System | SEC | unk | unk | unk | 0.04 | 0.5 | undet | unk | unk | Y | N (2015) |
| Common bottlenose dolphin | Northern Georgia/ Southern South Carolina Estuarine System | SEC | unk | unk | unk | 0.04 | 0.5 | unk | 1.4 | 1.4 | Y | N (2015) |
| Common bottlenose dolphin | Central Georgia Estuarine System | SEC | 192 | 0.04 | 185 | 0.04 | 0.5 | 1.9 | unk | unk | Y | N (2015) |
| Common bottlenose dolphin | Southern Georgia Estuarine System | SEC | 194 | 0.05 | 185 | 0.04 | 0.5 | 1.9 | unk | unk | Y | N (2015) |
| Common bottlenose dolphin | Jacksonville Estuarine System | SEC | unk | unk | unk | 0.04 | 0.5 | unk | 1.2 | 1.2 | Y | N (2015) |
| Common bottlenose dolphin | Indian River Lagoon Estuarine System | SEC | unk | unk | unk | 0.04 | 0.5 | unk | 4.4 | 4.4 | Y | N (2015) |
| Common bottlenose dolphin | Biscayne Bay | SEC | unk | unk | unk | 0.04 | 0.5 | unk | unk | unk | Y | N (2013) |
| Common bottlenose dolphin | Florida Bay | SEC | unk | unk | unk | 0.04 | 0.5 | undet | unk | unk | N | N (2013) |
| Harbor porpoise | Gulf of Maine/Bay of Fundy | NEC | 79,833 | 0.32 | 61,415 | 0.046 | 0.5 | 706 | 307 | 307 (0.16) | N | Y (m) |
| Harbor seal | Western North Atlantic | NEC | 75,834 | 0.15 | 66,884 | 0.12 | 0.5 | 2,006 | 368 | 356 (0.11) | N | Y (m) |

| Species | Stock Area | NMFS Ctr. | Nbest | Nbest CV | Nmin | Rmax | Fr | PBR | Total Annual S.I and Mort. | Annual Fish. S.I. and Mort. (cv) | Strategic Status | SAR Revised |
|------------------------------|--|-----------|-----------------------------|-------------|-----------------------------|------|------|-------------------------------|-------------------------------|-------------------------------------|---------------------|----------------|
| Gray seal | Western North Atlantic | NEC | 27,131 | 0.10 | 23,158 | 0.12 | 1.0 | 1,389 | 5,207 | 1,088 (0.09) | N | Y (a,m,p) |
| Harp seal | Western North Atlantic | NEC | unk | unk | unk | 0.12 | 1.0 | unk | 215,998 | 43 (0.24) | Ν | Y (2013) |
| Hooded seal | Western North Atlantic | NEC | unk | unk | unk | 0.12 | 0.75 | unk | 5,199 | 25(0.82) | N | N (2007) |
| Sperm whale | Gulf of Mexico | SEC | 763 | 0.38 | 560 | 0.04 | 0.1 | 1.1 | 0 | 0 | Y | N (2015) |
| Bryde's whale | Gulf of Mexico | SEC | 33 | 1.07 | 16 | 0.04 | 0.1 | 0.03 | 0.8 | 0 | Y | Y (m) |
| Cuvier's beaked whale | Gulf of Mexico | SEC | 74 | 1.04 | 36 | 0.04 | 0.5 | 0.4 | 0 | 0 | N | N (2012) |
| Blainville's beaked whale | Gulf of Mexico | SEC | 149 ^g | 0.91 | 77 | 0.04 | 0.5 | 0.8 | 0 | 0 | N | N (2012) |
| Gervais' beaked whale | Gulf of Mexico | SEC | 149 ^g | 0.91 | 77 | 0.04 | 0.5 | 0.8 | 0 | 0 | N | N (2012) |
| Common bottlenose dolphin | Gulf of Mexico, Continental shelf | SEC | 51,192 | 0.10 | 46,926 | 0.04 | 0.5 | 469 | 0.8 | 0.6 | N | N (2015) |
| Common bottlenose dolphin | Gulf of Mexico, eastern coastal | SEC | 12,388 | 0.13 | 11,110 | 0.04 | 0.5 | 111 | 1.6 | 1.6 | N | N (2015) |
| Common bottlenose dolphin | Gulf of Mexico, northern coastal | SEC | 7,185 | 0.21 | 6,044 | 0.04 | 0.5 | 60 | 0.4 | 0.4 | N | N (2015) |
| Common bottlenose dolphin | Gulf of Mexico, western coastal | SEC | 20,161 | 0.17 | 17,491 | 0.04 | 0.5 | 175 | 0.6 | 0.6 | N | N (2015) |
| Common bottlenose dolphin | Gulf of Mexico, Oceanic | SEC | 5,806 | 0.39 | 4,230 | 0.04 | 0.5 | 42 | 6.5 | 6.5 (0.65) | N | N (2014) |
| Common bottlenose dolphin | Gulf of Mexico, bay, sound and estuary (27 stocks) | SEC | unk for all but 2 stocks | unk | unk for all but 2 stocks | 0.04 | 0.5 | undet for all but 2 stocks | unk | unk | Y for all | N (2016) |

| Species | Stock Area | NMFS Ctr. | Nbest | Nbest CV | Nmin | Rmax | Fr | PBR | Total Annual S.I and Mort. | Annual Fish. S.I. and Mort. (cv) | Strategic Status | SAR Revised |
|------------------------------|--|-----------|--------|-------------|--------|------|-----|-------|-------------------------------|-------------------------------------|---------------------|----------------|
| Common bottlenose dolphin | Barataria Bay | SEC | 2,306 | 0.09 | 2,138 | 0.04 | 04 | 17 | 160 | 0.8 | Y | Y (a, m, p) |
| Common bottlenose dolphin | Mississippi Sound, Lake Borgne, Bay Boudreau | SEC | 3,046 | 0.06 | 2,896 | 0.04 | 0.4 | 23 | 310 | 1.0 | Y | Y(a, m, p) |
| Common bottlenose dolphin | St. Joseph Bay | SEC | 152 | 0.08 | unk | 0.04 | 0.5 | undet | unk | unk | Y | N (2015) |
| Common bottlenose dolphin | Choctawhatchee Bay | SEC | 179 | 0.04 | unk | 0.04 | 0.5 | undet | 0.4 | 0.4 | Y | N (2015) |
| Atlantic spotted dolphin | Gulf of Mexico | SEC | unk | unk | unk | 0.04 | 0.5 | undet | 42 | 42 (0.45) | N | N (2015) |
| Pantropical spotted dolphin | Gulf of Mexico | SEC | 50,880 | 0.27 | 40,699 | 0.04 | 0.5 | 407 | 4.4 | 4.4 | N | N (2015) |
| Striped dolphin | Gulf of Mexico | SEC | 1,849 | 0.77 | 1,041 | 0.04 | 0.5 | 10 | 0 | 0 | N | N (2012) |
| Spinner dolphin | Gulf of Mexico | SEC | 11,441 | 0.83 | 6,221 | 0.04 | 0.5 | 62 | 0 | 0 | N | N (2012) |
| Rough-toothed dolphin | Gulf of Mexico | SEC | 624 | 0.99 | 311 | 0.04 | 0.4 | 3 | 0.8 | 0.8 (1.0) | N | N (2016) |
| Clymene dolphin | Gulf of Mexico | SEC | 129 | 1.00 | 64 | 0.04 | 0.5 | 0.6 | 0 | 0 | N | N (2012) |
| Fraser's dolphin | Gulf of Mexico | SEC | unk | unk | unk | 0.04 | 0.5 | undet | 0 | 0 | N | N (2012) |
| Killer whale | Gulf of Mexico | SEC | 28 | 1.02 | 14 | 0.04 | 0.5 | 0.1 | 0 | 0 | N | N (2012) |
| False killer whale | Gulf of Mexico | SEC | unk | unk | unk | 0.04 | 0.5 | undet | 0 | 0 | N | N (2012) |
| Pygmy killer whale | Gulf of Mexico | SEC | 152 | 1.02 | 75 | 0.04 | 0.5 | 0.8 | 0 | 0 | N | N (2012) |

| Species | Stock Area | NMFS Ctr. | Nbest | Nbest CV | Nmin | Rmax | Fr | PBR | Total Annual S.I and Mort. | Annual Fish. S.I. and Mort. (cv) | Strategic Status | SAR Revised |
|--|--------------------------------------|-----------|-------------------|-------------|-------|------|-----|-----|-------------------------------|-------------------------------------|---------------------|----------------|
| Dwarf sperm whale | Gulf of Mexico | SEC | 186 ^h | 1.04 | 90 | 0.04 | 0.5 | 0.9 | 0 | 0 | N | N (2012) |
| Pygmy sperm whale | Gulf of Mexico | SEC | 186 ^h | 1.04 | 90 | 0.04 | 0.5 | 0.9 | 0.3 | 0.3 (1.0) | N | N (2012) |
| Melon-headed whale | Gulf of Mexico | SEC | 2,235 | 0.75 | 1,274 | 0.04 | 0.5 | 13 | 0 | 0 | N | N (2012) |
| Risso's dolphin | Gulf of Mexico | SEC | 2,442 | 0.57 | 1,563 | 0.04 | 0.5 | 16 | 7.9 | 7.9 (0.85) | N | N (2015) |
| Pilot whale, short- finned ⁱ | Gulf of Mexico | SEC | 2,415 | 0.66 | 1456 | 0.04 | 0.5 | 15 | 0.5 | 0.5 (1.0) | N | N (2015) |
| Sperm Whale | Puerto Rico and US Virgin Islands | SEC | unk | unk | unk | 0.04 | 0.1 | unk | unk | unk | Y | N (2010) |
| Common bottlenose dolphin | Puerto Rico and US Virgin Islands | SEC | unk | unk | unk | 0.04 | 0.5 | unk | unk | unk | Y | N (2011) |
| Cuvier's beaked whale | Puerto Rico and US Virgin Islands | SEC | unk | unk | unk | 0.04 | 0.5 | unk | unk | unk | Y | N (2011) |
| Pilot whale, short- finned | Puerto Rico and US Virgin Islands | SEC | unk | unk | unk | 0.04 | 0.5 | unk | unk | unk | Y | N (2011) |
| Spinner dolphin | Puerto Rico and US Virgin Islands | SEC | unk | unk | unk | 0.04 | 0.5 | unk | unk | unk | Y | N (2011) |
| Atlantic spotted dolphin | Puerto Rico and US Virgin Islands | SEC | unk | unk | unk | 0.04 | 0.5 | unk | unk | unk | Y | N (2011) |

a. The R given for right whales is the default Rmax of 0.04. The total estimated human-caused mortality and serious injury to right whales is estimated at 5.36 per year. This is derived from two components: 1) non-observed fishery entanglement records at 4.55 per year, and 2) ship strike records at 0.81 per year.

b. The total estimated human-caused mortality and serious injury to the Gulf of Maine humpback whale stock is estimated as 8.25 per year. This average is derived from two components: 1) incidental fishery interaction records 6.45; 2) records of vessel collisions, 1.8.

- c. The total estimated human-caused mortality and serious injury to the Western North Atlantic fin whale stock is estimated as 2.65 per year. This average is derived from two components: 1) incidental fishery interaction records 1.05; 2) records of vessel collisions, 1.6.
- d. The total estimated human-caused mortality and serious injury to the Nova Scotia sei whale stock is estimated as 0.8 per year. This average is derived from two components: 1) incidental fishery interaction records 0; 2) records of vessel collisions, 0.8.
- e. The total estimated human-caused mortality and serious injury to the Canadian East Coast minke whale stock is estimated as 9.0 per year. This average is derived from two components: 1) 7.6 minke whales per year (unknown CV) from U.S. and Canadian fisheries using strandings and entanglement data; and 3) 1.4 per year from vessel strikes.
- f. Estimates may include sightings of the coastal form.
- g. This estimate includes Gervais' beaked whales and Blainville's beaked whales for the Gulf of Mexico stocks, and all species of Mesoplodon in the Atlantic.
- h. This estimate includes both the dwarf and pygmy sperm whales.
- i. This estimate includes all *Globicephala sp.*, though it is presumed that only short-finned pilot whales are present in the Gulf of Mexico.